



Environment,
Land, Water
and Planning



Barwon Climate Projections 2019



Maximum and minimum daily temperatures will continue to increase over this century (very high confidence).



By the 2030s, increases in daily maximum temperature of 0.8 to 1.5°C (since the 1990s) are expected.



Rainfall will continue to be very variable over time, but over the long term it is expected to continue to decline in winter and spring (medium to high confidence) and autumn (low to medium confidence), but with some chance of little change.



Extreme rainfall events are expected to become more intense on average through the century (high confidence) but remain very variable in space and time.



By the 2050s, the climate of Geelong could be more like the current climate of Shepparton, and Colac more like Wodonga.

Introduction

Multiple lines of evidence indicate the global climate has changed predominantly due to human activity, and changes are projected to continue. Here we present projections of future climate change to help understand the scope and scale of changes to the climate we can expect locally in Barwon.

This report presents a snapshot of the *Victorian Climate Projections 2019* (VCP19) results for the Barwon region, and complements previous projections such as the *Victorian Climate Initiative*. Here we focus on the results from the new 5 km resolution downscaled climate simulations. Additional notes are provided drawing on earlier global climate model (GCM) and downscaled results where relevant.

Results are shown for two plausible scenarios of future greenhouse gas emissions: medium emissions (RCP4.5) and high emissions (RCP8.5). A more comprehensive set of results are provided in Tables 3–5 on the final pages. For more detail, please refer to the VCP19 Technical Report (TR).

Temperature

Victoria's temperature increased by just over 1°C between 1910 and 2018 (TR, Ch.5). This warming is expected to continue and, as a result, temperatures in the Barwon region will also increase. However, as Figure 1 shows, the amount of increase in the second half of the current century depends on the world's greenhouse gas emissions over the coming decades. Importantly, the upper range of temperature results from the VCP19 high-resolution modelling shows that a hotter future than that projected by the earlier GCM results is possible.

Under the high emissions scenario, maximum temperatures in the Barwon region are expected to show a median increase of 1.1°C by the 2030s (2020–2039), compared to 1986–2005. By mid-century, the increase is likely to be greater, with a median of 1.9°C. Under medium emissions, the mid-century maximum temperatures increase by a median of 1.4°C.

Increases in minimum temperatures are expected to be smaller, with a median of 0.7°C by the 2030s and 1.3°C by the 2050s (2040–2059) under high emissions.




Figure 1. Projected changes (compared to 1986-2005) in annual mean temperature for medium emissions (top) and high emissions (bottom). Light red bars: results from the full range of new downscaled modelling. Dark vertical line: median. Dark red bar: 2090s results from the most recent generation of global climate models for comparison.

Rainfall

Rainfall in Victoria has declined in most seasons over recent decades, with the greatest decreases in the cooler seasons (TR, Ch.5). Barwon’s rainfall is naturally highly variable and this natural variability will dominate the rainfall over the next decade or so.

Over time, annual rainfall totals are likely to decline, particularly under high emissions (see Figure 2), with the greatest drying in spring. By late-century under high emissions, the climate change trend becomes obvious compared to natural variability (not shown) with a median of 24% decrease in annual totals, larger (34%) in spring.




Figure 2. Projected changes (compared to 1986-2005) in annual mean rainfall for medium emissions (top) and high emissions (bottom). Bars show the 10th to 90th percentile range. Blue bars: results from the new downscaled modelling. Dark vertical line: median. Dark blue dots: individual models. Green bar: results from all available modelling (high resolution and GCM) for comparison at 2090.

Extremes

As warming continues, we expect more heat extremes in the decades to come. Projections show that in the 2050s under high emissions, the 1-in-20-year hottest summer day is likely to increase by a median value of 2.7°C compared to 1986–2005 (see Figure 3). Under medium emissions, the median increase for the 2050s is 1.8°C.




Figure 3. Projected changes in the 1-in-20-year hottest day for 2040–2059 compared to 1986–2005. Blue bars (left): medium emissions. Purple bars (right): high emissions. Dark horizontal lines: median.

Another way to look at extreme temperatures is to look at the number of temperature exceedances per year. For example, on average between 1981 and 2010, Geelong experienced 6.4 days per year when the temperature exceeded 35°C (see Table 1). By the 2050s under high emissions, this is expected to increase to between 9 and 16 days. The increase is slightly less under medium emissions, reaching 8 to 13 days on average. Importantly for many people, minimum (usually overnight) temperatures are also expected to increase. For example, Geelong’s daily minimum temperature exceeded 20°C on average 3.9 days per year between 1981 and 2010. Under high emissions, by the 2050s, these hot nights are expected to occur 7.6 to 12.2 days per year on average.

Table 1. Historic and projected days (median, 10th and 90th percentile) per year with maximum temperature greater than 35°C for two locations in 2040–2059. Projected values for medium emissions (RCP4.5) are in blue (top); high emissions (RCP8.5) are in red (bottom).

Town	Days/year above 35°C	
	1981–2010	2040–2059
Geelong	6.4	RCP4.5 10.9 (7.7 to 12.6)
		RCP8.5 12.6 (9.1 to 15.6)
Colac	7.4	RCP4.5 12.9 (8.9 to 14.0)
		RCP8.5 14.4 (11.1 to 20.6)

A warmer climate is expected to bring more heavy rainfall events, but variability in high rainfall events is naturally large so there is a large range of possibilities in any 20-year period. VCP19 shows a range of median changes in annual extreme daily rainfall for a 20-year return period between -4 to +10% by the 2050s for the Barwon region, depending on the season and emissions scenario (see Figure 4).




Figure 4. Projected percentage change in the 1-in-20-year maximum daily rainfall for 2040–2059 compared to the 1986–2005. Blue bars (left): medium emissions. Purple bars (right): high emissions. Dark horizontal lines: median.

Frost

In a warming climate, frosts are expected to become less frequent over time. However, it is possible for there to be an increased risk of frost in some regions and seasons when cold clear nights persist longer than is suggested by the projected change in minimum temperature (TR, Ch.5). Over time the effect of increasing minimum temperatures is expected to gradually overpower the other effects and lead to a decrease in frost risk in almost all regions and seasons.




Figure 5. Projected changes in the 1-in-20-year coldest daily minimum temperature for 2040–2059 compared to 1986–2005. Blue bars (left): medium emissions. Purple bars (right): high emissions. Dark horizontal lines: median.

Keeping this in mind, it is useful to look at projected changes in very cold days. By the 2050s under high emissions, the 1-in-20-year coldest minimum winter daily temperature is projected to warm by a median of 0.8°C. Under medium emissions, the median increase in the coldest daily minimum temperature is almost 0.5°C (see Figure 5).

As with hot days, it is useful to look at minimum temperature thresholds (Table 2). Historically from 1981 to 2010, Geelong experienced on average 3.8 days per year below 2°C whereas Colac experienced 5.8 days. By the 2050s under high emissions, this may decline to between 0.7 and 1.5 days per year for Geelong and 0.8 to 3.0 days per year for Colac. Under medium emissions, the expected reductions are less.

Table 2. Historic and projected days (median, 10th and 90th percentile) per year with minimum temperature less than 2°C for two locations in 2040–2059. Projected values for medium emissions (RCP4.5) are in blue (top); high emissions (RCP8.5) are in red (bottom).

Town	Days/year below 2°C	
	1981–2010	2040–2059
Geelong	3.8	RCP4.5 1.5 (0.9 to 2.9)
		RCP8.5 1.0 (0.7 to 1.5)

Colac	5.8	RCP4.5 2.4 (1.7 to 3.5)
		RCP8.5 1.5 (0.8 to 3.0)

Fire danger

Fire weather has become more dangerous in southern Australia since the mid-1900s. In addition, fire seasons have become longer and commenced earlier over this time. This is likely to continue, with the severity of ongoing changes dependent on the emissions pathway that the world follows.

For example, there is high confidence that the number of fire days where the Forest Fire Danger Index is greater than the 95th percentile for 1986–2005 is projected to increase at Geelong by a median value of 9 days per year by the 2050s under high emissions (or a 49% increase).

Sea level

Sea levels around the Australian coastline rose at an average rate of 2.1 mm/yr between 1966 and 2009. From 1993 to 2009, this increased to 3.1 mm/yr. Sea levels are expected to continue rising through the 21st century (very high confidence). For example, at Geelong by the 2050s under high emissions, sea level is expected to rise by a median value of 24 cm (4 mm/yr) compared to the 1990s (TR, Ch.5).



Mark Chew

Barwon Climate Projections 2019

Table 3. Summary of changes (compared to 1986-2005) for selected climate variables (median, 10th and 90th percentile) from new 5 km downscaled results for greenhouse gas emissions scenarios RCP 4.5 (blue, 1st row) and RCP 8.5 (red, 2nd row). Bold black values (3rd row) for rainfall represent the projected range of change in rainfall using a distribution based on all climate modelling available, past and present.

Variable	Season	1986–2005 average Geelong ¹	Emissions scenario	Projected changes		
				2020–2039	2040–2059	2080–2099
Maximum temperature (°C)	Annual	19.1°C	RCP4.5	0.9 (0.8 to 1.2)	1.4 (0.9 to 1.9)	2.1 (1.7 to 3.1)
			RCP8.5	1.1 (0.9 to 1.5)	1.9 (1.4 to 2.6)	3.8 (2.6 to 4.9)
	Summer	23.8°C	RCP4.5	0.9 (0.6 to 1.8)	1.5 (0.9 to 2.5)	2.2 (1.7 to 3.2)
			RCP8.5	1.0 (0.7 to 1.9)	1.8 (1.4 to 3.0)	3.7 (2.1 to 5.3)
	Autumn	19.9°C	RCP4.5	0.9 (0.5 to 1.2)	1.3 (0.9 to 1.8)	1.9 (1.7 to 3.2)
			RCP8.5	1.1 (0.8 to 1.4)	1.6 (1.5 to 2.5)	3.8 (2.7 to 4.9)
	Winter	14.5°C	RCP4.5	0.7 (0.6 to 0.8)	1.1 (0.7 to 1.2)	1.7 (1.3 to 2.5)
			RCP8.5	0.8 (0.6 to 1.0)	1.5 (1.3 to 1.7)	3.3 (2.5 to 3.9)
	Spring	18.3°C	RCP4.5	1.3 (1.0 to 1.5)	1.8 (1.2 to 2.3)	2.7 (2.0 to 3.4)
			RCP8.5	1.3 (1.1 to 1.9)	2.2 (1.6 to 3.2)	4.5 (2.9 to 5.7)
Minimum temperature (°C)	Annual	9.0°C	RCP4.5	0.6 (0.6 to 0.7)	1.1 (0.8 to 1.4)	1.7 (1.2 to 2.0)
			RCP8.5	0.7 (0.6 to 0.9)	1.3 (1.1 to 1.8)	2.7 (2.3 to 3.6)
	Summer	12.5°C	RCP4.5	0.9 (0.5 to 1.0)	1.2 (0.9 to 2.0)	2.0 (1.1 to 2.2)
			RCP8.5	0.8 (0.6 to 1.3)	1.6 (1.3 to 2.1)	3.1 (2.5 to 4.2)
	Autumn	9.7°C	RCP4.5	0.6 (0.4 to 0.8)	1.1 (0.7 to 1.4)	1.6 (1.1 to 2.2)
			RCP8.5	0.7 (0.4 to 0.9)	1.3 (0.9 to 1.9)	2.9 (2.4 to 3.9)
	Winter	5.9°C	RCP4.5	0.6 (0.3 to 0.8)	0.9 (0.6 to 1.2)	1.4 (1.1 to 2.2)
			RCP8.5	0.6 (0.5 to 0.8)	1.1 (1.0 to 1.3)	2.6 (2.2 to 3.1)
	Spring	7.9°C	RCP4.5	0.6 (0.4 to 0.8)	0.9 (0.7 to 1.4)	1.6 (1.1 to 1.7)
			RCP8.5	0.8 (0.5 to 1.0)	1.2 (1.1 to 1.8)	2.5 (1.9 to 3.2)
Rainfall (%)	Annual	548.4 mm	RCP4.5	-7 (-14 to -1)	-6 (-12 to -3)	-10 (-21 to 1)
			RCP8.5	-10 (-19 to -2)	-11 (-20 to -4)	(-27 to +5)
	Summer	114.2 mm	RCP4.5	-5 (-12 to 9)	-1 (-5 to 11)	-16 (-38 to 25)
			RCP8.5	-10 (-22 to 18)	-9 (-18 to 2)	(-34 to +17)
	Autumn	132.9 mm	RCP4.5	-5 (-22 to 2)	-9 (-17 to 11)	-9 (-27 to 0)
			RCP8.5	-12 (-18 to -1)	-10 (-18 to 1)	(-24 to +7)
	Winter	141.2 mm	RCP4.5	-9 (-12 to -6)	-7 (-10 to -4)	-7 (-12 to -1)
			RCP8.5	-10 (-16 to 0)	-11 (-18 to -7)	(-40 to -1)
	Spring	159.1 mm	RCP4.5	-10 (-19 to 10)	-12 (-22 to 2)	-21 (-26 to -5)
			RCP8.5	-18 (-25 to -1)	-22 (-26 to 15)	(-29 to +18)

Notes:

1. Bureau of Meteorology monthly station data from Grovedale (Geelong Airport) (1986–2005).

Barwon Climate Projections 2019

Table 4. Summary of changes (compared to 1986-2005) for selected climate variables (median, 10th and 90th percentile) from new 5 km downscaled results for greenhouse gas emissions scenarios RCP 4.5 (blue, 1st row) and RCP 8.5 (red, 2nd row).

Variable	Season	Emissions scenario	Projected changes		
			2020–2039	2040–2059	2080–2099
Relative humidity (%)	Annual	RCP4.5	-1.2 (-2.2 to -0.1)	-1.0 (-2.0 to -0.1)	-1.9 (-3.9 to -0.5)
		RCP8.5	-1.4 (-2.6 to -0.3)	-1.7 (-2.9 to -0.1)	-3.9 (-5.2 to 1.0)
	Summer	RCP4.5	-0.9 (-3.5 to -0.3)	-1.0 (-2.5 to 0.1)	-2.8 (-4.5 to 0.0)
		RCP8.5	-0.9 (-4.1 to -0.1)	-2.0 (-3.7 to -0.1)	-2.8 (-5.0 to 2.9)
	Autumn	RCP4.5	-1.4 (-2.4 to 0.8)	-0.8 (-2.0 to 0.3)	-1.4 (-3.7 to 0.4)
		RCP8.5	-1.3 (-2.7 to -0.2)	-1.9 (-3.0 to 0.2)	-4.4 (-4.8 to 1.9)
	Winter	RCP4.5	-0.3 (-0.8 to 0.0)	-0.3 (-1.0 to 0.2)	-0.7 (-1.9 to -0.1)
		RCP8.5	-0.4 (-0.9 to -0.1)	-0.7 (-1.4 to -0.2)	-2.1 (-3.2 to -0.3)
	Spring	RCP4.5	-2.0 (-2.8 to -0.8)	-2.0 (-3.7 to -0.9)	-3.6 (-5.5 to -1.5)
		RCP8.5	-2.5 (-3.3 to -0.9)	-3.3 (-4.7 to -0.1)	-6.2 (-8.1 to -0.7)
Pan evaporation (%)	Annual	RCP4.5	7.6 (6.3 to 12.2)	11.1 (5.6 to 15.5)	16.2 (12.8 to 25.7)
		RCP8.5	8.5 (6.0 to 11.8)	14.6 (8.0 to 21.5)	29.5 (12.5 to 43.2)
	Summer	RCP4.5	11.4 (5.3 to 24.1)	14.7 (8.5 to 30.4)	25.9 (19.0 to 36.6)
		RCP8.5	9.9 (7.9 to 22.2)	19.5 (13.8 to 36.8)	43.4 (14.5 to 64.4)
	Autumn	RCP4.5	6.6 (-0.4 to 8.2)	7.0 (1.8 to 9.6)	10.9 (6.4 to 18.3)
		RCP8.5	6.9 (2.7 to 9.9)	10.5 (4.2 to 12.1)	21.8 (7.2 to 31.7)
	Winter	RCP4.5	1.5 (1.2 to 4.8)	2.4 (-0.1 to 6.9)	6.1 (3.0 to 12.0)
		RCP8.5	2.2 (0.5 to 4.9)	4.4 (2.6 to 7.1)	11.2 (5.1 to 19.0)
	Spring	RCP4.5	13.4 (10.9 to 15.6)	17.7 (12.0 to 20.5)	27.4 (18.1 to 36.6)
		RCP8.5	13.4 (9.7 to 15.5)	22.1 (10.9 to 32.7)	44.8 (22.8 to 58.3)
Solar radiation (%)	Annual	RCP4.5	1.8 (0.8 to 3.5)	2.2 (1.4 to 3.9)	3.2 (2.4 to 6.5)
		RCP8.5	2.0 (1.2 to 2.9)	3.3 (1.3 to 4.5)	6.2 (1.4 to 7.7)
	Summer	RCP4.5	2.0 (-0.9 to 3.1)	1.5 (-0.8 to 3.2)	2.5 (-0.8 to 4.1)
		RCP8.5	0.6 (-1.2 to 1.9)	1.4 (-0.4 to 3.6)	2.7 (-2.0 to 4.1)
	Autumn	RCP4.5	0.7 (-0.8 to 4.7)	2.4 (-0.6 to 3.9)	2.1 (0.7 to 6.1)
		RCP8.5	3.1 (0.7 to 4.1)	3.0 (0.3 to 4.7)	6.8 (1.0 to 8.7)
	Winter	RCP4.5	1.9 (1.3 to 2.8)	3.0 (0.4 to 3.7)	3.7 (1.8 to 7.3)
		RCP8.5	2.5 (1.2 to 3.4)	3.5 (3.0 to 5.9)	9.2 (3.8 to 9.7)
	Spring	RCP4.5	3.5 (2.8 to 3.8)	4.5 (3.3 to 6.1)	6.9 (4.9 to 9.9)
		RCP8.5	3.7 (2.5 to 4.5)	5.5 (2.3 to 7.8)	9.7 (5.3 to 11.7)
Surface wind speed (%)	Annual	RCP4.5	-1.0 (-2.3 to 1.1)	-1.7 (-2.7 to 1.0)	-1.2 (-2.0 to 0.0)
		RCP8.5	-2.2 (-2.9 to 0.8)	-2.6 (-3.3 to -0.3)	-2.5 (-5.5 to 0.2)
	Summer	RCP4.5	0.3 (-2.0 to 1.5)	-0.5 (-3.0 to 0.5)	-0.3 (-0.9 to 0.4)
		RCP8.5	0.4 (-1.4 to 1.6)	-0.4 (-2.5 to 1.1)	-0.6 (-3.1 to 0.9)
	Autumn	RCP4.5	-0.4 (-4.3 to 0.3)	-2.6 (-4.3 to 2.4)	-2.8 (-5.6 to -0.4)
		RCP8.5	-2.0 (-4.5 to 0.0)	-3.5 (-4.1 to 0.5)	-4.8 (-6.8 to -0.8)
	Winter	RCP4.5	-2.7 (-4.9 to 3.8)	-2.2 (-5.3 to 3.3)	-0.3 (-3.2 to 2.7)
		RCP8.5	-3.7 (-4.8 to 3.1)	-3.5 (-5.8 to 1.1)	-3.6 (-9.7 to 1.5)
	Spring	RCP4.5	0.0 (-1.9 to 2.0)	-0.8 (-2.4 to 1.7)	-0.6 (-1.7 to 0.6)
		RCP8.5	-1.7 (-4.6 to 0.6)	-2.1 (-3.4 to 0.2)	-2.0 (-5.0 to 2.3)

Barwon Climate Projections 2019

Table 5. Summary of changes (compared to 1986-2005) for selected climate extremes (median, 10th and 90th percentile) from new 5 km downscaled results for greenhouse gas emissions scenarios RCP 4.5 (blue, 1st row) and RCP 8.5 (red, 2nd row).

Variable	Season	Emissions scenario	Projected changes		
			2020–2039	2040–2059	2080–2099
Extreme (ARI20) daily maximum temperature (°C)	Annual	RCP4.5	0.4 (-0.4 to 2.4)	1.6 (-0.9 to 4.0)	3.0 (1.6 to 4.3)
		RCP8.5	0.2 (-0.3 to 2.7)	2.6 (1.5 to 5.2)	4.4 (2.2 to 5.8)
	Summer	RCP4.5	0.4 (-0.3 to 2.7)	1.8 (-1.2 to 4.0)	3.4 (1.3 to 4.3)
		RCP8.5	0.3 (-0.4 to 3.1)	2.7 (1.1 to 5.3)	4.5 (2.0 to 5.9)
	Autumn	RCP4.5	2.3 (0.6 to 4.2)	1.3 (-0.8 to 4.0)	4.8 (1.5 to 5.9)
		RCP8.5	3.0 (-1.7 to 4.2)	4.0 (2.0 to 4.7)	4.1 (3.4 to 6.9)
	Winter	RCP4.5	1.6 (0.9 to 2.3)	1.6 (0.2 to 4.3)	2.9 (1.6 to 5.6)
		RCP8.5	1.9 (-0.7 to 3.2)	2.9 (0.2 to 4.3)	6.5 (2.9 to 10.5)
	Spring	RCP4.5	1.5 (-0.3 to 3.0)	2.5 (1.0 to 3.7)	4.0 (1.8 to 4.4)
		RCP8.5	1.6 (-0.3 to 3.1)	3.2 (1.0 to 4.5)	5.4 (3.9 to 7.4)
Extreme (ARI20) daily rainfall (%)	Annual	RCP4.5	-4 (-19 to 29)	7 (4 to 21)	-1 (-11 to 37)
		RCP8.5	2 (-20 to 17)	-2 (-10 to 24)	15 (-5 to 34)
	Summer	RCP4.5	-2 (-19 to 49)	10 (-1 to 32)	-6 (-23 to 28)
		RCP8.5	1 (-30 to 21)	0 (-10 to 10)	13 (-4 to 23)
	Autumn	RCP4.5	0 (-20 to 19)	2 (-11 to 28)	-2 (-12 to 40)
		RCP8.5	-6 (-26 to 18)	8 (-18 to 42)	20 (-25 to 49)
	Winter	RCP4.5	-3 (-19 to 31)	4 (-17 to 33)	1 (-12 to 24)
		RCP8.5	-2 (-18 to 18)	4 (-13 to 15)	20 (-12 to 25)
	Spring	RCP4.5	2 (-24 to 19)	-4 (-13 to 13)	6 (-20 to 31)
		RCP8.5	-3 (-27 to 32)	6 (-13 to 17)	-13 (-27 to 19)

© 2019 State of Victoria

Creative Commons Attribution 4.0 Australian licence (<https://creativecommons.org/licenses/by/4.0/>)

Updated 19th February 2020

ISBN 978-1-76077-733-3 (Print)

ISBN 978-1-76077-734-0 (pdf/online/MS word)

Citation: Clarke JM, Grose M, Thatcher M, Round V & Heady C. 2019. Barwon Climate Projections 2019. CSIRO, Melbourne Australia.

This report should be read in conjunction with Clarke JM, Grose M, Thatcher M, Hernaman V, Heady C, Round V, Rafter T, Trenham C & Wilson L. 2019. Victorian Climate Projections 2019 Technical Report. CSIRO, Melbourne Australia.

Disclaimer: CSIRO and the State of Victoria advise that the information contained in this publication comprises general statements based on scientific research. The reader is advised and needs to be aware that such information may be incomplete or unable to be used in any specific situation. No reliance or actions must therefore be made on that information without seeking prior expert professional, scientific and technical advice. To the extent permitted by law, CSIRO and the State of Victoria (including their employees and consultants) exclude all liability to any person for any consequences, including but not limited to all losses, damages, costs, expenses and any other compensation, arising directly or indirectly from using this publication (in part or in whole) and any information or material contained in it.